

Addendum to: The Infrared Spectrum of CS₂ in the ν₃ Band Region

F. Winther

Institut für Physikalische Chemie, Universität Kiel

Z. Naturforsch. **44a**, 594 (1989);
received April 1, 1989

In a recent paper by Wells et al. [1] the band origins of ν₃ of ¹²C³²S₂ and ¹³C³²S₂ have been determined with high accuracy from heterodyne frequency measurements. As mentioned in our paper on these bands [2] our spectra were calibrated with water vapour lines from [3]. This limited the accuracy of all band centers in the paper to $\pm 3 \cdot 10^{-4} \text{ cm}^{-1}$ (one standard error) – not allowing for any systematic error of the calibration lines, however.

Reprint requests to Dr. F. Winther, Institut für Physikalische Chemie der Christian-Albrechts-Universität zu Kiel, Ludwig-Meyn-Str. 8, D-2300 Kiel.

- [1] J. S. Wells, M. Schneider, and A. G. Maki, *J. Mol. Spectr.* **132**, 422 (1988).
 [2] F. Winther, U. Heyne, and A. Guarnieri, *Z. Naturforsch.* **43a**, 215 (1988).
 [3] G. Guelachvili, *J. Opt. Soc. Amer.* **73**, 137 (1983).

In a paper by Brown and Toth [4] it has been indicated that the frequencies of these calibration lines are $3.63 \cdot 10^{-4} \text{ cm}^{-1}$ too high (mean value for 71 lines). It should be noted that a corresponding correction has not been applied to the same frequencies in “Handbook of Infrared Standards” [5].

Indeed, a comparison of the band origins of [1] and [2] shows that our frequencies are 2.9(8) and $5.2(8) \cdot 10^{-4} \text{ cm}^{-1}$ too high for ¹²C³²S₂ and ¹³C³²S₂, respectively. This confirms the statement of [4] above about the water line frequencies.

In our spectra ¹³C³²S₂ was recorded in natural abundance so that the differences from the heterodyne measurements refer to the strongest and one of the weakest line series. There may have been a slight asymmetry in our line profiles (which we did not check), which would account for the incomplete numerical agreement for the two bands. However, a mean correction of $4 \cdot 10^{-4} \text{ cm}^{-1}$ should be subtracted from all band center frequencies given in our paper [2]. Their absolute accuracies should then all be about 10^{-4} cm^{-1} .

- [4] L. R. Brown and R. A. Toth, *J. Opt. Soc. Amer.* **B2**, 842 (1985).
 [5] G. Guelachvili and K. N. Rao, *Handbook of Infrared Standards*, Academic Press, NY 1986

0932-0784 / 89 / 0600-0594 \$ 01.30/0. – Please order a reprint rather than making your own copy.

